# antibodies - online.com











### Overview

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | AMH  |
| Binding Specificity: | AA 19-560  |
| Reactivity:          | Human  |
| Host:                | Mouse  |
| Clonality:           | Monoclonal   |
| Conjugate:           | This AMH antibody is un-conjugated   |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

### **Product Details**

| Purpose:          | Monoclonal Antibody to Anti-Mullerian Hormone (AMH)   |
|-------------------|---|
| Immunogen:        | Recombinant Anti-Mullerian Hormone (AMH) corresdonding to Leu19~Arg560 with N-terminal His Tag  |
| Clone:            | H13   |
| Isotype:          | IgG1 kappa  |
| Specificity:      | The antibody is a mouse monoclonal antibody raised against AMH. It has been selected for its ability to recognize AMH in immunohistochemical staining and western blotting. |
| Cross-Reactivity: | Rat   |
| Purification:     | Protein A + Protein G affinity chromatography   |

## **Target Details**

Expiry Date:


24 months

| Larget Details      |  |
|---------------------|--|
| Target:             | AMH  |
| Alternative Name:   | Anti-Mullerian Hormone (AMH Products)  |
| Background:         | MIF, MIH, MIS, Müllerian Inhibiting Factor, Müllerian Inhibiting Hormone, Müllerian Inhibiting   |
|                     | Substance  |
| Pathways:           | Negative Regulation of Hormone Secretion   |
| Application Details |  |
| Application Notes:  | Western blotting: 0.5-2 μg/mL  |
|                     | 1:500-2000 Immunohistochemistry: 5-20 μg/mL  |
|                     | 1:50-200 Immunocytochemistry: 5-20 µg/mL   |
|                     | 1:50-200 Optimal working dilutions must be determined by end user.                               |
| Comment:            | The thermal stability is described by the loss rate. The loss rate was determined by accelerated |
|                     | thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious          |
|                     | degradation and precipitation were observed. The loss rate is less than 5% within the expiration |
|                     | date under appropriate storage condition.  |
| Restrictions:       | For Research Use only  |
| Handling            |  |
| Format:             | Liquid   |
| Buffer:             | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.                                      |
| Preservative:       | Sodium azide   |
| Precaution of Use:  | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which                    |
|                     | should be handled by trained staff only.   |
| Storage:            | 4 °C,-20 °C  |
| Storage Comment:    | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without  |
|                     | detectable loss of activity. Avoid repeated freeze-thaw cycles.                                  |



# **Western Blotting**

**Image 1.** Detection of AMH in Rat Testis lysate using Monoclonal Antibody to Anti-Mullerian Hormone (AMH)



#### **Immunohistochemistry**

**Image 2.** Detection of AMH in Human Hela cell using Monoclonal Antibody to Anti-Mullerian Hormone (AMH)