
antibodies - online.com







anti-EGF antibody (AA 977-1029)





()	1 /	\sim	KI /	110	Νę
	1//	\vdash	I \/	1 ←	٠// ٢

Quantity:	100 μL	
Target:	EGF	
Binding Specificity:	AA 977-1029	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This EGF antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

Product Details

Purpose:	Polyclonal Antibody to Epidermal Growth Factor (EGF)	
Immunogen:	Recombinant Epidermal Growth Factor (EGF) corresdonding to Asn977~Arg1029 with N-terminal His and GST Tag	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against EGF. It has been selected for its ability to recognize EGF in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Human, Rat	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	

Target Details


Target:	EGF	
Alternative Name:	Epidermal Growth Factor (EGF Products)	
Background:	URG, Beta-Urogastrone, Pro-epidermal growth factor	
Pathways:	NF-kappaB Signaling, RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Regulation of Carbohydrate Metabolic Process, Hepatitis C, Protein targeting to Nucleus, Interaction of EGFR with phospholipase C-gamma, Thromboxane A2 Receptor Signaling, EGFR Downregulation	

Application Details

Application Notes:	Western blotting: 0.5-2 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-		
	20 μg/mL Optimal working dilutions must be determined by end user.		
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated		
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious		
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration		
	date under appropriate storage condition.		
Restrictions:	For Research Use only		


Handling

riarianing		
Format:	Liquid	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	24 months	



Immunohistochemistry

Image 1. #VALUE!



Western Blotting

Image 2. Detection of Recombinant EGF, Mouse using Polyclonal Antibody to Epidermal Growth Factor (EGF)