
antibodies - online.com







anti-TGFB1 antibody (AA 279-390)



Images



\sim				
	$ V \cap$	r\/I	19	٨

Quantity:	100 μL
Target:	TGFB1
Binding Specificity:	AA 279-390
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TGFB1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Transforming Growth Factor Beta 1 (TGFb1)	
Immunogen:	Recombinant Transforming Growth Factor Beta 1 (TGFb1) corresdonding to Ala279~Ser390 with N-terminal His Tag	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against TGFb1. It has been selected for its ability to recognize TGFb1 in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Human, Mouse	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	

Target Details

Alternative Name: Tran	nsforming Growth Factor Beta 1 (TGFB1 Products)
Background: TGF	-B1, CED, DPD1, LAP, Camurati-Engelmann Disease, Latency-associated peptide
Baci Imm Prod Res Reg	R Signaling Pathway, Dopaminergic Neurogenesis, Cellular Response to Molecule of terial Origin, Glycosaminoglycan Metabolic Process, Regulation of Leukocyte Mediated nunity, Regulation of Muscle Cell Differentiation, Positive Regulation of Immune Effector cess, Cell-Cell Junction Organization, Production of Molecular Mediator of Immune conse, Ribonucleoside Biosynthetic Process, Skeletal Muscle Fiber Development, ulation of Carbohydrate Metabolic Process, Protein targeting to Nucleus, Autophagy, cer Immune Checkpoints

Application Details		
Application Notes:	Western blotting: 0.5-2 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-20 μg/mL Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	

detectable loss of activity. Avoid repeated freeze-thaw cycles.

Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without

4 °C,-20 °C


Storage:

Storage Comment:

Expiry Date:


24 months

Images




Immunohistochemistry

Image 1. Detection of TGFb1 in Rat Cardiac Muscle Tissue using Polyclonal Antibody to Transforming Growth Factor Beta 1 (TGFb1)



Western Blotting

Image 2. Detection of Recombinant TGFb1, Rat using Polyclonal Antibody to Transforming Growth Factor Beta 1 (TGFb1)



Immunofluorescence

Image 3. Detection of TGFb1 in Human U2OS cell using Polyclonal Antibody to Transforming Growth Factor Beta 1 (TGFb1)

Please check the product details page for more images. Overall 8 images are available for ABIN7429469.