



[Go to Product page](#)

Datasheet for ABIN7429550
anti-CST3 antibody (AA 21-140)

3 Images

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | CST3 |
| Binding Specificity: | AA 21-140 |
| Reactivity: | Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CST3 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

Product Details

| | |
|-------------------|--|
| Purpose: | Polyclonal Antibody to Cystatin C (Cys-C) |
| Immunogen: | Recombinant Cystatin C (Cys-C) corresponding to Ala21~Ala140 with N-terminal His Tag |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against Cys-C. It has been selected for its ability to recognize Cys-C in immunohistochemical staining and western blotting. |
| Cross-Reactivity: | Rat |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography |

Target Details


| | |
|-------------------|--|
| Target: | CST3 |
| Alternative Name: | Cystatin C (CST3 Products) |
| Background: | CST3, Cystatin 3, Gamma Trace, Post-Gamma-Globulin, Neuroendocrine Basic Polypeptide, Amyloid Angiopathy And Cerebral Hemorrhage |

Application Details

| | |
|--------------------|---|
| Application Notes: | Western blotting: 0.5-2 µg/mL Immunohistochemistry: 5-20 µg/mL Immunocytochemistry: 5-20 µg/mL Optimal working dilutions must be determined by end user. |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions: | For Research Use only |


Handling

| | |
|--------------------|---|
| Format: | Liquid |
| Buffer: | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles. |
| Expiry Date: | 24 months |




Immunohistochemistry

Image 1. Detection of Cys-C in Mouse Cerebrum Tissue using Polyclonal Antibody to Cystatin C (Cys-C)



Western Blotting

Image 2. Detection of Recombinant CST3, Mouse using Polyclonal Antibody to Cystatin C (Cys-C)



Immunohistochemistry

Image 3. Detection of Cys-C in Mouse Kidney Tissue using Polyclonal Antibody to Cystatin C (Cys-C)