

Datasheet for ABIN7429733 anti-EPO antibody (AA 41-206)

1 Image



Overview

| Quantity: | 100 μL |
|----------------------|--|
| Target: | EPO |
| Binding Specificity: | AA 41-206 |
| Reactivity: | Dog |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This EPO antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

Product Details

| Purpose: | Polyclonal Antibody to Erythropoietin (EPO) |
|-------------------|--|
| Immunogen: | Recombinant Erythropoietin (EPO) corresdonding to Ala41~Arg206 with N-terminal His Tag |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against EPO. It has been selected for its ability to recognize EPO in immunohistochemical staining and western blotting. |
| Cross-Reactivity: | Goat, Sheep |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography |

Target Details

| Target: | EPO EPO |
|-------------------|---|
| Alternative Name: | Erythropoietin (EPO Products) |
| Target Type: | Hormone |
| Background: | EP, Epoetin, Erythropoetin, Hematopoietin, Hemopoietin |
| UniProt: | P33707 |
| Pathways: | JAK-STAT Signaling, Hormone Activity, Negative Regulation of intrinsic apoptotic Signaling, Negative Regulation of Transporter Activity |

Application Details

| Application Notes: | Western blotting: 0.5-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-20 μg/mL,Optimal working dilutions must be determined by end user. |
|--------------------|--|
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated |
| | thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration |
| Restrictions: | date under appropriate storage condition. For Research Use only |

Handling

| Format: | Liquid |
|--------------------|---|
| Concentration: | 500 μg/mL |
| Buffer: | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles. |
| Expiry Date: | 24 months |



Western Blotting

Image 1. Detection of EPO in Caprine Kidney lysate using Polyclonal Antibody to Erythropoietin (EPO)