antibodies .- online.com







anti-IL-8 antibody (AA 28-99)





Overview

Quantity:	100 μL
Target:	IL-8 (IL8)
Binding Specificity:	AA 28-99
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IL-8 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Interleukin 8 (IL8)
Immunogen:	Recombinant Interleukin 8 (IL8) corresdonding to Ser28~Ser99 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against IL8. It has been selected for its ability to recognize IL8 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	IL-8 (IL8)

Target Details

Alternative Name:	Interleukin 8 (IL8 Products)
Background:	CXCL8, AMCF-I, GCP1, K60, LECT, LUCT, LYNAP, MDNCF, MONAP, NAF, NAP1, SCYB8, TSG1 B-ENAP, Neutrophil-Activating Protein 1, Granulocyte Chemotactic Protein 1
Pathways:	TLR Signaling, Cellular Response to Molecule of Bacterial Origin, Regulation of G-Protein Coupled Receptor Protein Signaling, ER-Nucleus Signaling, Hepatitis C, Autophagy
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL

Restrictions:	For Research Use only
	date under appropriate storage condition.
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	Optimal working dilutions must be determined by end user.
	Immunocytochemistry: 5-20 μg/mL
	Immunohistochemistry: 5-20 μg/mL
, ipplication i totoo.	. 160.6


Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months




Western Blotting

Image 1. Detection of IL8 in Human Serum using Polyclonal Antibody to Interleukin 8 (IL8)



Western Blotting

Image 2. Detection of Recombinant IL8, Human using Polyclonal Antibody to Interleukin 8 (IL8)



Western Blotting

Image 3. Detection of IL8 in Human Leukocyte lysate using Polyclonal Antibody to Interleukin 8 (IL8)