# antibodies -online.com





## anti-CXCL1 antibody (AA 35-107)

2 Images



Go to Product page

#### Overview

Quantity:	100 μL
Target:	CXCL1
Binding Specificity:	AA 35-107
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CXCL1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunohistochemistry (Paraffinembedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**


Purpose:	Polyclonal Antibody to Chemokine (C-X-C Motif) Ligand 1 (CXCL1)
Immunogen:	Recombinant Chemokine (C-X-C Motif) Ligand 1 (CXCL1) corresdonding to Ala35~Asn107 with N-terminal His and GST Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CXCL1. It has been selected for its ability to recognize CXCL1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Pig
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

### **Target Details**

Expiry Date:


24 months

rarget Details	
Target:	CXCL1
Alternative Name:	Chemokine (C-X-C Motif) Ligand 1 (CXCL1 Products)
Background:	NAP3, GRO1, GROa, GRO-A, MGSA-A, NAP3, SCYB1, Chemokine C-X-C-Motif Ligand 1, Melanoma Growth Stimulating Activity Alpha, Fibroblast Secretory Protein
Pathways:	Autophagy
Application Details	
Application Notes:	Western blotting: 1-5 $\mu$ g/mL Immunocytochemistry in formalin fixed cells: 5-20 $\mu$ g/mL Immunohistochemistry in formalin fixed frozen section: 5-20 $\mu$ g/mL Immunohistochemistry in paraffin section: 5-20 $\mu$ g/mL Enzyme-linked Immunosorbent Assay: 0.05-2 $\mu$ g/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.



#### **Immunohistochemistry**

**Image 1.** Detection of CXCL1 in Human Cerebrum Tissue using Polyclonal Antibody to Chemokine (C-X-C Motif) Ligand 1 (CXCL1)



#### **Western Blotting**

Image 2. Detection of Recombinant NAP3, Human using Polyclonal Antibody to Chemokine (C-X-C Motif) Ligand 1 (CXCL1)