antibodies -online.com





anti-CXCL1 antibody (AA 25-96)

2 Images



Go to Product page

Overview

Quantity:	100 μL
Target:	CXCL1
Binding Specificity:	AA 25-96
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CXCL1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Chemokine (C-X-C Motif) Ligand 1 (CXCL1)
Immunogen:	Recombinant Chemokine (C-X-C Motif) Ligand 1 (CXCL1) corresdonding to Ala25~Lys96 (Accession # P14095)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CXCL1. It has been selected for its ability to recognize CXCL1 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target: CXCL1


Target Details

Expiry Date:

Target Details	
Alternative Name:	Chemokine (C-X-C Motif) Ligand 1 (CXCL1 Products)
Background:	NAP3, GRO1, GROa, GRO-A, MGSA-A, NAP3, SCYB1, Chemokine C-X-C-Motif Ligand 1,
	Melanoma Growth Stimulating Activity Alpha, Fibroblast Secretory Protein
Pathways:	Autophagy
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL
	1:250-1000 Immunohistochemistry: 5-20 μg/mL
	1:25-100 Immunocytochemistry: 5-20 µg/mL
	1:25-100 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without


detectable loss of activity. Avoid repeated freeze-thaw cycles.

24 months



Immunohistochemistry

Image 1. Detection of CXCL1 in Rat Testis Tissue using Polyclonal Antibody to Chemokine (C-X-C Motif) Ligand 1 (CXCL1)



Western Blotting

Image 2. Detection of Recombinant NAP3, Rat using Polyclonal Antibody to Chemokine (C-X-C Motif) Ligand 1 (CXCL1)