

Datasheet for ABIN7430564 anti-HGF antibody (AA 495-728)

1 Image



Go to Product page

_				
()	VA	rv	IPI	٨

Quantity:	100 μL
Target:	HGF
Binding Specificity:	AA 495-728
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HGF antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)
	Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Hepatocyte Growth Factor (HGF)	
Immunogen:	Recombinant Hepatocyte Growth Factor (HGF) corresdonding to Val495~Ser728 with N-terminal His Tag	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against HGF. It has been selected for its ability to recognize HGF in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Mouse, Pig, Rat	
Purification:	ication: Antigen-specific affinity chromatography followed by Protein A affinity chromatography	

Target Details

Alternative Name: Hepatocyte	Hepatocyte Growth Factor (HGF Products)	
	HGF, F-TCF, HGFB, HPTA, SF, Scatter Factor, Hepapoietin A, Fibroblast-Derived Tumor Cytotoxic Factor, Lung Fibroblast-Derived Mitogen	
,	ng, Carbohydrate Homeostasis, Glycosaminoglycan Metabolic Process, Synaptic Signaling of Hepatocyte Growth Factor Receptor	

Application Details


Application Notes:

	Immunohistochemistry: 5-20 μg/mL
	Immunocytochemistry: 5-20 μg/mL
	Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only

Western blotting: $0.5-2 \mu g/mL$

Handling

Format:	Liquid	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	24 months	



Western Blotting

Image 1. Detection of Recombinant HGF, Human using Polyclonal Antibody to Hepatocyte Growth Factor (HGF)