
antibodies -online.com









Go to Product page

| \sim | | |
|-----------|----------|--------------------------|
| () , , , | V / / | 1014 |
| 1 11// | ۱ / ۱ عد | $1 \leftarrow 1 \land 1$ |
| | | |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | TIMP2 |
| Binding Specificity: | AA 27-220 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

Product Details

| Purpose: | Polyclonal Antibody to Tissue Inhibitors Of Metalloproteinase 2 (TIMP2) |
|-------------------|--|
| Immunogen: | Recombinant Tissue Inhibitors Of Metalloproteinase 2 (TIMP2) corresdonding to Cys27~Pro220 with N-terminal His Tag |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against TIMP2. It has been selected for its ability to recognize TIMP2 in immunohistochemical staining and western blotting. |
| Cross-Reactivity: | Mouse, Rat |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography |

Target Details

|--|

Target Details


| Alternative Name: | Tissue Inhibitors Of Metalloproteinase 2 (TIMP2 Products) |
|-------------------|---|
| Background: | CSC-21K, TIMP Metallopeptidase Inhibitor 2 |
| Pathways: | cAMP Metabolic Process |

Application Details

| Application Notes: | Western blotting: 0.5-2 μg/mL |
|--------------------|--|
| | Immunohistochemistry: 5-20 μg/mL |
| | Immunocytochemistry: 5-20 μg/mL |
| | Optimal working dilutions must be determined by end user. |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated |
| | thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious |
| | degradation and precipitation were observed. The loss rate is less than 5% within the expiration |
| | date under appropriate storage condition. |
| Restrictions: | For Research Use only |


Handling

| Format: | Liquid |
|--------------------|---|
| Buffer: | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles. |
| Expiry Date: | 24 months |




Immunohistochemistry

Image 1. Detection of TIMP2 in Human Kidney Tissue using
Polyclonal Antibody to Tissue Inhibitors Of
Metalloproteinase 2 (TIMP2)



Western Blotting

Image 2. Detection of Recombinant TIMP2, Human using Polyclonal Antibody to Tissue Inhibitors Of Metalloproteinase 2 (TIMP2)



Immunohistochemistry

Image 3. Detection of TIMP2 in Human Mammary gland Tissue using Polyclonal Antibody to Tissue Inhibitors Of Metalloproteinase 2 (TIMP2)

Please check the product details page for more images. Overall 4 images are available for ABIN7431134.