

Datasheet for ABIN743189

anti-IKBKG antibody (pSer31) (Cy7)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	IKBKG
Binding Specificity:	pSer31
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IKBKG antibody is conjugated to Cy7
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human IKK gamma around the phosphorylation site of Ser31
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified by Protein A.

Target Details

Target:	IKBKG
Alternative Name:	IKK gamma (IKBKG Products)

Target Details

Background:	Synonyms: Ikb kinase associated protein 1, Ikb kinase subunit gamma, Inhibitor of nuclear factor kappa B kinase subunit gamma, AMCBX1, FIP 3, FIP3, Fip3p, I kappa B kinase gamma, Ikb kinase associated protein 1, Ikb kinase gamma subunit, Ikb kinase subunit gamma, IKBKG, IKKAP 1, IKKAP1, IKKG, IKKgamma,IKK gamma. Background: Pro inflammatory cytokines activate the transcription factor NF kappa B by stimulating the activity of a protein kinase that phosphorylates I kappa B, an inhibitor of NF kappa B, at sites that trigger its ubiquitination and degradation. A large, cytokine responsive I kappa B kinase (IKK) complex has been purified and the genes encoding 2 of its subunits have been cloned. These subunits, IKK alpha and Ikk beta, are protein kinases whose function is needed for NF kappa B activation by pro inflammatory stimuli. IKK is composed of similar amounts of IKK alpha, Ikk beta, which are differentially processed forms of a third subunit, IKK gamma. IKK gamma interacts preferentially with IKK beta and is required for the activation of the IKK complex.
Gene ID:	8517
Pathways:	NF-kappaB Signaling , RTK Signaling , TCR Signaling , TLR Signaling , Fc-epsilon Receptor Signaling Pathway , Activation of Innate immune Response , M Phase , Production of Molecular Mediator of Immune Response , Hepatitis C , Protein targeting to Nucleus , Toll-Like Receptors Cascades , BCR Signaling , Ubiquitin Proteasome Pathway , S100 Proteins

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months