# antibodies -online.com





## anti-Tissue factor antibody (AA 30-251)

2 Images



#### Go to Product page

_					
U	V	er	V	Ie	W

Quantity:	100 μL
Target:	Tissue factor (F3)
Binding Specificity:	AA 30-251
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Tissue factor antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

### **Product Details**

Purpose:	Polyclonal Antibody to Tissue Factor (TF)
Immunogen:	Recombinant Tissue Factor (TF) corresdonding to Gly30~Glu251 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TF. It has been selected for its ability to recognize TF in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

Target: Tissue factor (F3)

## **Target Details**


Alternative Name:	Tissue Factor (F3 Products)
Background:	CD142, FIII, F3, TFA, Thromboplastin, Coagulation Factor III
Pathways:	Positive Regulation of Endopeptidase Activity, Smooth Muscle Cell Migration, Platelet-derived growth Factor Receptor Signaling

## **Application Details**

Application Notes:	Western blotting: 0.5-2 μg/mL
	Immunohistochemistry: 5-20 μg/mL
	Immunocytochemistry: 5-20 μg/mL
	Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only


## Handling

Format:	Liquid	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	24 months	



### **Immunohistochemistry**

**Image 1.** Detection of TF in Mouse Pancreas Tissue using Polyclonal Antibody to Tissue Factor (TF)



### **Western Blotting**

Image 2. Detection of Recombinant TF, Mouse using Polyclonal Antibody to Tissue Factor (TF)