# antibodies -online.com





## anti-ICAM-3/CD50 antibody (AA 259-491)

2 Images



#### Go to Product page

$\sim$			
/ I\	/arv	/1/	711

Quantity:	100 μL	
Target:	ICAM-3/CD50 (ICAM3)	
Binding Specificity:	AA 259-491	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ICAM-3/CD50 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	
Draduat Dataila		

#### **Product Details**

Purpose:	Polyclonal Antibody to Intercellular Adhesion Molecule 3 (ICAM3)
Immunogen:	Recombinant Intercellular Adhesion Molecule 3 (ICAM3) corresdonding to Gly259~Val491 (Accession # P32942)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against ICAM3. It has been selected for its ability to recognize ICAM3 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

#### **Target Details**

Target: ICAM-3/CD50 (ICAM3)


### **Target Details**

Alternative Name:	Intercellular Adhesion Molecule 3 (ICAM3 Products)	
Background:	CDW50, CD50, ICAM-R	
Application Details		
Application Notes:	Western blotting: 0.5-2 μg/mL	
	1:150-700 Immunohistochemistry: 5-20 μg/mL	
	1:15-70 Immunocytochemistry: 5-20 μg/mL	
	1:15-70 Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be	
	handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	
	detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	24 months	



#### **Immunohistochemistry**

Image 1. #VALUE!



#### **Western Blotting**

Image 2. Detection of Recombinant ICAM3, Human using Polyclonal Antibody to Intercellular Adhesion Molecule 3 (ICAM3)