

Datasheet for ABIN7432001 anti-TIMP1 antibody (AA 24-178)

Image

Target:



Go	to	Product	page

Overview		
Quantity:	100 μL	
Target:	TIMP1	
Binding Specificity:	AA 24-178	
Reactivity:	Horse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TIMP1 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	
Product Details		
Purpose:	Polyclonal Antibody to Tissue Inhibitors Of Metalloproteinase 1 (TIMP1)	
Immunogen:	Recombinant Tissue Inhibitors Of Metalloproteinase 1 (TIMP1) corresdonding to Cys24~Ser178 with N-terminal His and GST Tag	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against TIMP1. It has been selected for its ability to recognize TIMP1 in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	


TIMP1

Target Details

Expiry Date:

24 months

rarget Details		
Alternative Name:	TIMP1 (TIMP1 Products)	
Background:	EPO, CLGI, EPA, HCI, Erythroid Potentiating Activity, Collagenase Inhibitor, Metalloproteinase	
	inhibitor 1, Fibroblast collagenase inhibitor	
UniProt:	002722	
Application Details		
Application Notes:	Western blotting: 0.5-2 μg/mLlmmunohistochemistry: 5-20 μg/mLlmmunocytochemistry: 5-20	
	μg/mLOptimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	500 μg/mL	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be	
	handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	
	detectable loss of activity. Avoid repeated freeze-thaw cycles.	



Western Blotting

Image 1. Detection of Recombinant TIMP1, Equine using Polyclonal Antibody to Tissue Inhibitors Of Metalloproteinase 1 (TIMP1)