# antibodies -online.com







## anti-CXCL5 antibody (AA 45-118)





Go to Product page

| $\sim$ |           |      |    |   |
|--------|-----------|------|----|---|
|        | $ V \cap$ | r\/I | 19 | ٨ |

| Quantity:            | 100 μL   |  |
|----------------------|--|--|
| Target:              | CXCL5  |  |
| Binding Specificity: | AA 45-118  |  |
| Reactivity:          | Mouse  |  |
| Host:                | Rabbit   |  |
| Clonality:           | Polyclonal   |  |
| Conjugate:           | This CXCL5 antibody is un-conjugated   |  |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |  |

### **Product Details**


| Purpose:      | Polyclonal Antibody to Epithelial Neutrophil Activating Peptide 78 (ENA78)   |  |
|---------------|--|--|
| Immunogen:    | Recombinant Epithelial Neutrophil Activating Peptide 78 (ENA78) corresdonding to Val45~Ala118 (Accession # P50228)   |  |
| Isotype:      | IgG  |  |
| Specificity:  | The antibody is a rabbit polyclonal antibody raised against ENA78. It has been selected for its ability to recognize ENA78 in immunohistochemical staining and western blotting. |  |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography   |  |

## Target Details

|--|


## **Target Details**

| rarget betails      |  |  |
|---------------------|--|--|
| Alternative Name:   | Epithelial Neutrophil Activating Peptide 78 (CXCL5 Products)                                     |  |
| Background:         | CXCL5, SCYB5, Chemokine C-X-C-Motif Ligand 5, Small Inducible Cytokine Subfamily B(Cys-X-        |  |
|                     | Cys),Member 5, Neutrophil-activating peptide ENA-78  |  |
| Pathways:           | Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity     |  |
| Application Details |  |  |
| Application Notes:  | Western blotting: 0.5-2 μg/mL  |  |
|                     | Immunohistochemistry: 5-20 μg/mL   |  |
|                     | Immunocytochemistry: 5-20 μg/mL  |  |
|                     | Optimal working dilutions must be determined by end user.  |  |
| Comment:            | The thermal stability is described by the loss rate. The loss rate was determined by accelerated |  |
|                     | thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious          |  |
|                     | degradation and precipitation were observed. The loss rate is less than 5% within the expiration |  |
|                     | date under appropriate storage condition.  |  |
| Restrictions:       | For Research Use only  |  |
| Handling            |  |  |
| Format:             | Liquid   |  |
| Buffer:             | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.                                 |  |
| Preservative:       | ProClin  |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be               |  |
|                     | handled by trained staff only.   |  |
| Storage:            | 4 °C,-20 °C  |  |
| Storage Comment:    | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without  |  |
|                     | detectable loss of activity. Avoid repeated freeze-thaw cycles.                                  |  |
| Expiry Date:        | 24 months  |  |
|                     |  |  |



### **Immunohistochemistry**

Image 1. #VALUE!



### **Western Blotting**

**Image 2.** Detection of Recombinant ENA78, Mouse using Polyclonal Antibody to Epithelial Neutrophil Activating Peptide 78 (ENA78)