
antibodies -online.com







Images



Overview

Quantity:	100 μL
Target:	SFTPD
Binding Specificity:	AA 21-375
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SFTPD antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Surfactant Associated Protein D (SPD)
Immunogen:	Recombinant Surfactant Associated Protein D (SPD) corresdonding to Ala21~Phe375 linked with AEMKTYSPEF with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against SPD. It has been selected for its ability to recognize SPD in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography


Target Details

Expiry Date:

Target Details	
Target:	SFTPD
Alternative Name:	Surfactant Associated Protein D (SFTPD Products)
Background:	SFTPD, COLEC7, PSP-D, SFTP4, SP-D, Pulmonary Surfactant Protein D, Collectin-7, Lung surfactant protein D
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-20 μg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without


detectable loss of activity. Avoid repeated freeze-thaw cycles.

24 months




Immunohistochemistry

Image 1. Detection of SPD in Human Lung Tissue using Polyclonal Antibody to Surfactant Associated Protein D (SPD)



Western Blotting

Image 2. Detection of SPD in Human Lung Tissue using Polyclonal Antibody to Surfactant Associated Protein D (SPD)



Western Blotting

Image 3. Detection of Recombinant SPD, Human using Polyclonal Antibody to Surfactant Associated Protein D (SPD)