
antibodies -online.com







anti-Elastin antibody (AA 392-645)



Images



| () | 11 | \sim | rv | | ۱ ۸ |
|-----|-------|--------|------|---|-----|
| | 1 \ / | ⊢ | I \/ | ╙ | 1/1 |
| | | | | | |

| Quantity: | 100 μL |
|----------------------|---|
| Target: | Elastin (ELN) |
| Binding Specificity: | AA 392-645 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Elastin antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunohistochemistry (Paraffinembedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

Product Details


| Purpose: | Polyclonal Antibody to Elastin (ELN) |
|-------------------|--|
| Immunogen: | Recombinant Elastin (ELN) corresdonding to Gly392~Ala645 with N-terminal His Tag |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against ELN. It has been selected for its ability to recognize ELN in immunohistochemical staining and western blotting. |
| Cross-Reactivity: | Pig, Rat |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography |

Target Details

| Target: | Elastin (ELN) |
|---------------------|--|
| Alternative Name: | Elastin (ELN Products) |
| Background: | WBS, WS, SVAS, Tropoelastin, Supravalvular Aortic Stenosis, Williams-Beuren Syndrome |
| Pathways: | Regulation of Actin Filament Polymerization |
| | |
| Application Details | |


| Application Details | |
|---------------------|---|
| Application Notes: | Western blotting: 0.5-2 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-20 μg/mL Optimal working dilutions must be determined by end user. |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. |
| Preservative: | ProClin |

| Format: | Liquid | |
|--------------------|---|--|
| Buffer: | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | |
| Storage: | 4 °C,-20 °C | |
| Storage Comment: | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 24 months | |




Immunohistochemistry

Image 1. Detection of ELN in Human Spleen Tissue using Polyclonal Antibody to Elastin (ELN)



Western Blotting

Image 2. Detection of ELN in Human Lung Tissue using Polyclonal Antibody to Elastin (ELN)



Western Blotting

Image 3. Detection of Recombinant ELN, Human using Polyclonal Antibody to Elastin (ELN)

Please check the product details page for more images. Overall 8 images are available for ABIN7433489.