
# antibodies - online.com







# anti-ADCY1 antibody (AA 835-1061)

**Images** 



#### Overview

Quantity:	100 μL
Target:	ADCY1
Binding Specificity:	AA 835-1061
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ADCY1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

## **Product Details**

Purpose:	Polyclonal Antibody to Adenylate Cyclase 1, Brain (ADCY1)
Immunogen:	Recombinant Adenylate Cyclase 1, Brain (ADCY1) corresdonding to Arg835~Asn1061 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against ADCY1. It has been selected for its ability to recognize ADCY1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography


# **Target Details**

cyclase type I  Pathways: EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Thyroid Hormone Synthesis	get Details	
Background:  AC1, ATP pyrophosphate-lyase 1, Ca(2+)/calmodulin-activated adenylyl cyclase, Adeny cyclase type I  Pathways:  EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Thyroid Hormone Synthesis Metabolic Process, Myometrial Relaxation and Contraction, G-protein mediated Events Interaction of EGFR with phospholipase C-gamma	jet:	ADCY1
cyclase type I  Pathways: EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Thyroid Hormone Synthesis  Metabolic Process, Myometrial Relaxation and Contraction, G-protein mediated Events  Interaction of EGFR with phospholipase C-gamma	rnative Name:	Adenylate Cyclase 1, Brain (ADCY1 Products)
Metabolic Process, Myometrial Relaxation and Contraction, G-protein mediated Events Interaction of EGFR with phospholipase C-gamma	kground:	AC1, ATP pyrophosphate-lyase 1, Ca(2+)/calmodulin-activated adenylyl cyclase, Adenylate cyclase type I
Application Details	ıways:	EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Thyroid Hormone Synthesis, cAMP Metabolic Process, Myometrial Relaxation and Contraction, G-protein mediated Events, Interaction of EGFR with phospholipase C-gamma
	olication Details	
Application Notes: Western blotting: 0.5-2 µg/mL Immunohistochemistry: 5-20 µg/mL Immunocytochemi	lication Notes:	Western blotting: 0.5-2 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-

Application Notes:	Western blotting: 0.5-2 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-
	20 μg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only


# Handling

Tarianing	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



## **Immunohistochemistry**

Image 1. #VALUE!



## **Western Blotting**

Image 2. Detection of Recombinant ADCY1, Human using Polyclonal Antibody to Adenylate Cyclase 1, Brain (ADCY1)