# antibodies -online.com





## anti-Dystrophin antibody (AA 3059-3314)



Image



#### Overview

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | Dystrophin (DMD)   |
| Binding Specificity: | AA 3059-3314   |
| Reactivity:          | Mouse  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This Dystrophin antibody is un-conjugated  |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

#### **Product Details**

| Purpose:          | Polyclonal Antibody to Dystrophin (DMD)  |
|-------------------|--|
| Immunogen:        | Recombinant Dystrophin (DMD) corresdonding to Ser3059~Ile3314 with N-terminal His Tag  |
| Isotype:          | IgG  |
| Specificity:      | The antibody is a rabbit polyclonal antibody raised against DMD. It has been selected for its ability to recognize DMD in immunohistochemical staining and western blotting. |
| Cross-Reactivity: | Human, Rat   |
| Purification:     | Antigen-specific affinity chromatography followed by Protein A affinity chromatography   |

### **Target Details**

Precaution of Use:

Storage Comment:

Storage:

Expiry Date:

| Target:           | Dystrophin (DMD)                                     |
|-------------------|--|
| Alternative Name: | Dystrophin (DMD Products)                            |
| Background:       | CMD3B, Muscular Dystrophy, Duchenne And Becker Types |
| Pathways:         | Skeletal Muscle Fiber Development                    |

| Application Details |   |
|---------------------|---|
| Application Notes:  | Western blotting: 0.5-2 μg/mL<br>1:500-2000 Immunohistochemistry: 5-20 μg/mL  |
|                     | 1:50-200 Immunocytochemistry: 5-20 µg/mL<br>1:50-200 Optimal working dilutions must be determined by end user.  |
| Comment:            | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Liquid  |
| Buffer:             | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.   |
| Preservative:       | Sodium azide  |

should be handled by trained staff only.


4 °C,-20 °C

24 months

This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without

detectable loss of activity. Avoid repeated freeze-thaw cycles.



#### **Western Blotting**

Image 1. Detection of Recombinant DMD, Mouse using Polyclonal Antibody to Dystrophin (DMD)