

Datasheet for ABIN7433768
anti-IDO antibody (AA 2-403)

5 Images

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	IDO
Binding Specificity:	AA 2-403
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IDO antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Indoleamine-2,3-Dioxygenase (IDO)
Immunogen:	Recombinant Indoleamine-2,3-Dioxygenase (IDO) corresponding to Ala2~Gly403 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against IDO. It has been selected for its ability to recognize IDO in immunohistochemical staining and western blotting.
Cross-Reactivity:	Pig, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details


Target:	IDO
Alternative Name:	Indoleamine-2,3-Dioxygenase (IDO Products)
Background:	CD107B, INDO, Indoleamine-Pyrrole 2,3 Dioxygenase

Application Details

Application Notes:	Western blotting: 1-5 µg/mL Immunocytochemistry in formalin fixed cells: 5-20 µg/mL Immunohistochemistry in formalin fixed frozen section: 5-20 µg/mL Immunohistochemistry in paraffin section: 5-20 µg/mL Enzyme-linked Immunosorbent Assay: 0.05-2 µg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months




Western Blotting

Image 1. Detection of IDO in Human Placenta lysate using Polyclonal Antibody to Indoleamine-2,3-Dioxygenase (IDO)



Western Blotting

Image 2. Detection of Recombinant IDO, Human using Polyclonal Antibody to Indoleamine-2,3-Dioxygenase (IDO)



Immunohistochemistry

Image 3. Detection of IDO in Human Kidney Tissue using Polyclonal Antibody to Indoleamine-2,3-Dioxygenase (IDO)

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN7433768.