# antibodies -online.com





## anti-F13B antibody (AA 21-208)





#### Overview


| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | F13B   |
| Binding Specificity: | AA 21-208  |
| Reactivity:          | Mouse  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This F13B antibody is un-conjugated  |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

#### **Product Details**

| Purpose:          | Polyclonal Antibody to Coagulation Factor XIII B Polypeptide (F13B)  |
|-------------------|--|
| Immunogen:        | Recombinant Coagulation Factor XIII B Polypeptide (F13B) corresdonding to Ala21~Gln208 with N-terminal His Tag   |
| Isotype:          | IgG  |
| Specificity:      | The antibody is a rabbit polyclonal antibody raised against F13B. It has been selected for its ability to recognize F13B in immunohistochemical staining and western blotting. |
| Cross-Reactivity: | Rat  |
| Purification:     | Antigen-specific affinity chromatography followed by Protein A affinity chromatography   |

### **Target Details**

| Target:             | F13B   |
|---------------------|--|
| Alternative Name:   | Coagulation Factor XIII B Polypeptide (F13B Products)  |
| Background:         | F13-B, FXIIIB, Fibrin-stabilizing factor B subunit, Protein-glutamine gamma-glutamyltransferase  |
|                     | B chain, Transglutaminase B chain  |
| Application Details |  |
| Application Notes:  | Western blotting: 0.5-2 μg/mL  |
|                     | Immunohistochemistry: 5-20 μg/mL   |
|                     | Immunocytochemistry: 5-20 μg/mL  |
|                     | Optimal working dilutions must be determined by end user.  |
| Comment:            | The thermal stability is described by the loss rate. The loss rate was determined by accelerated |
|                     | thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious          |
|                     | degradation and precipitation were observed. The loss rate is less than 5% within the expiration |
|                     | date under appropriate storage condition.  |
| Restrictions:       | For Research Use only  |
| Handling            |  |
| Format:             | Liquid   |
| Buffer:             | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.                                      |
| Preservative:       | Sodium azide   |
| Precaution of Use:  | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which                    |
|                     | should be handled by trained staff only.   |
| Storage:            | 4 °C,-20 °C  |
| Storage Comment:    | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without  |
|                     | detectable loss of activity. Avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:        | 24 months  |



#### **Western Blotting**

**Image 1.** Detection of Recombinant F13B, Mouse using Polyclonal Antibody to Coagulation Factor XIII B Polypeptide (F13B)