

Datasheet for ABIN7434434  
**anti-C8G antibody (AA 39-202)**

## 3 Images

[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	C8G
Binding Specificity:	AA 39-202
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This C8G antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

## Product Details

Purpose:	Polyclonal Antibody to Complement Component 8g (C8g)
Immunogen:	Recombinant Complement Component 8g (C8g) corresponding to Lys39~Arg202 with N-terminal His and GST Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against C8g. It has been selected for its ability to recognize C8g in immunohistochemical staining and western blotting.
Cross-Reactivity:	Pig
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

## Target Details


Target:	C8G
Alternative Name:	Complement Component 8g ( <a href="#">C8G Products</a> )
Pathways:	<a href="#">Complement System</a>

## Application Details

Application Notes:	Western blotting: 0.5-2 µg/mL Immunohistochemistry: 5-20 µg/mL Immunocytochemistry: 5-20 µg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



#### Immunohistochemistry

**Image 1.** Detection of C8g in Human Kidney Tissue using Polyclonal Antibody to Complement Component 8g (C8g)



#### Western Blotting

**Image 2.** Detection of C8g in Human Serum using Polyclonal Antibody to Complement Component 8g (C8g)

#### Western Blotting

**Image 3.** Detection of Recombinant C8g, Human using Polyclonal Antibody to Complement Component 8g (C8g)