
antibodies -online.com







anti-Complement Factor I antibody (AA 340-583)



Image



\sim	
()\/\Di	view
	VICVV

Quantity:	100 μL
Target:	Complement Factor I (CFI)
Binding Specificity:	AA 340-583
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Complement Factor I antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Complement Factor I (CFI)
Immunogen:	Recombinant Complement Factor I (CFI) corresdonding to Ile340~Val583 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CFI. It has been selected for its ability to recognize CFI in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target: Complement Factor I (CFI)

Target Details

Storage Comment:


Expiry Date:

Target Details	
Alternative Name:	Complement Factor I (CFI Products)
Background:	CF-I, C3b-INA, FI, IF, KAF, Factor I, Konglutinogen-Activating Factor, C3b-Inactivator, C3B/C4B inactivator
Pathways:	Complement System
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-20 μg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C

detectable loss of activity. Avoid repeated freeze-thaw cycles.

24 months

Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without



Western Blotting

Image 1. Detection of Recombinant CFI, Human using Polyclonal Antibody to Complement Factor I (CFI)