antibodies -online.com







anti-HMGA1 antibody





Overview

Quantity:	100 μL
Target:	HMGA1
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HMGA1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details


Target:

Purpose:	Polyclonal Antibody to High Mobility Group AT Hook Protein 1 (HMGA1)
Immunogen:	Recombinant High Mobility Group AT Hook Protein 1 (HMGA1)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against HMGA1. It has been selected for its ability to recognize HMGA1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

HMGA1

Target Details

rarget Details	
Alternative Name:	High Mobility Group AT Hook Protein 1 (HMGA1 Products)
Background:	HMG-R, HMGIY, High mobility group protein A1, High mobility group protein R, High mobility
	group protein HMG-I/HMG-Y
Pathways:	Nuclear Hormone Receptor Binding, Autophagy
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-
	$20~\mu g/mL$ Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without
	detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



Western Blotting

Image 1. Detection of Recombinant HMGA1, Human using Polyclonal Antibody to High Mobility Group AT Hook Protein 1 (HMGA1)