



[Go to Product page](#)

Datasheet for ABIN7434617
anti-HMGA2 antibody (AA 1-109)

1 Image

Overview

Quantity:	100 µL
Target:	HMGA2
Binding Specificity:	AA 1-109
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HMGA2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunocytochemistry (ICC), Immunohistochemistry (IHC), Immunoprecipitation (IP)

Product Details

Purpose:	Polyclonal Antibody to High Mobility Group AT Hook Protein 2 (HMGA2)
Immunogen:	Recombinant High Mobility Group AT Hook Protein 2 (HMGA2) corresponding to Met1~Asp109 with N-terminal His and GST Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against HMGA2. It has been selected for its ability to recognize HMGA2 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	HMGA2
---------	-------

Target Details


Alternative Name:	High Mobility Group AT Hook Protein 2 (HMGA2 Products)
Background:	BABL, HMGI-C, HMGIC, LIPO, High mobility group protein HMGI-C
Pathways:	Positive Regulation of Response to DNA Damage Stimulus

Application Details

Application Notes:	Western blotting: 0.5-2 µg/mL 1:500-2000 Immunohistochemistry: 5-20 µg/mL 1:50-200 Immunocytochemistry: 5-20 µg/mL 1:50-200 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



Western Blotting

Image 1. Detection of Recombinant HMGA2, Human using Polyclonal Antibody to High Mobility Group AT Hook Protein 2 (HMGA2)