antibodies -online.com





anti-Dystroglycan antibody (AA 28-406)

3 Images



Overview

Quantity:	100 μL
Target:	Dystroglycan (DAG1)
Binding Specificity:	AA 28-406
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Dystroglycan antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details


Purpose:	Polyclonal Antibody to Dystrophin Associated Glycoprotein 1 (DAG1)
Immunogen:	Recombinant Dystrophin Associated Glycoprotein 1 (DAG1) corresdonding to His28~Pro406 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against DAG1. It has been selected for its ability to recognize DAG1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Human, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	Dystroglycan (DAG1)
Alternative Name:	Dystrophin Associated Glycoprotein 1 (DAG1 Products)
Background:	Dystroglycan 1, 156DAG, A3a, AGRNR, DAG, Alpha-Dystroglycan, Beta-Dystroglycan
Pathways:	Maintenance of Protein Location, Regulation of Carbohydrate Metabolic Process, Protein targeting to Nucleus


Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-20 μg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months




Western Blotting

Image 1. Detection of DAG1 in Mouse Placenta lysate using Polyclonal Antibody to Dystrophin Associated Glycoprotein 1 (DAG1)



Western Blotting

Image 2. Detection of Recombinant DAG1, Mouse using Polyclonal Antibody to Dystrophin Associated Glycoprotein 1 (DAG1)



Western Blotting

Image 3. Detection of DAG1 in Mouse Skeletal muscle lysate using Polyclonal Antibody to Dystrophin Associated Glycoprotein 1 (DAG1)