antibodies .- online.com







anti-Urocortin 2 antibody (AA 23-107)



Images



_					
U	V	er	V	Ie	W

Quantity:	100 μL
Target:	Urocortin 2 (UCN2)
Binding Specificity:	AA 23-107
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Urocortin 2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Target:

Purpose:	Polyclonal Antibody to Urocortin 2 (UCN2)	
Immunogen:	Recombinant Urocortin 2 (UCN2) corresdonding to Pro23~Gly107 (Accession # Q91WW1)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against UCN2. It has been selected for its ability to recognize UCN2 in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	
Target Details		

Urocortin 2 (UCN2)

Target Details


Alternative Name:	Urocortin 2 (UCN2 Products)
Background:	URP, SRP, UCN-II, UCNII, Stresscopin-Related Peptide, Urocortin-related peptide
Pathways:	Positive Regulation of Peptide Hormone Secretion, Negative Regulation of Hormone Secretion, cAMP Metabolic Process

Application Details

Restrictions:	For Research Use only		
	date under appropriate storage condition.		
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration		
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious		
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated		
	Optimal working dilutions must be determined by end user.		
	Immunocytochemistry: 5-20 μg/mL		
	Immunohistochemistry: 5-20 μg/mL		
Application Notes:	Western blotting: 0.5-2 μg/mL		


Handling

Format:	Liquid
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months




Immunohistochemistry

Image 1. Detection of UCN2 in Human Testis Tissue using Polyclonal Antibody to Urocortin 2 (UCN2)



Western Blotting

Image 2. Detection of Recombinant UCN2, Rat using Polyclonal Antibody to Urocortin 2 (UCN2)



Immunohistochemistry

Image 3. Detection of UCN2 in Human Tonsil Tissue using Polyclonal Antibody to Urocortin 2 (UCN2)