
antibodies - online.com







anti-Oxytocin Receptor antibody (AA 43-129)



Image



$\overline{}$				
\bigcap	10	r\/I	101	Λ

Quantity:	100 μL
Target:	Oxytocin Receptor (OXTR)
Binding Specificity:	AA 43-129
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Oxytocin Receptor antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Oxytocin Receptor (OXTR)	
Immunogen:	Recombinant Oxytocin Receptor (OXTR) corresdonding to Val43~Leu129 with N-terminal His and GST Tag	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against OXTR. It has been selected for its ability to recognize OXTR in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	

Target Details

or (OXTR)	Oxytocin Receptor (OXTR)	Target:
-----------	--------------------------	---------

Target Details


Alternative Name:	Oxytocin Receptor (OXTR Products)	
Background:	OT-R	
Pathways:	Myometrial Relaxation and Contraction, Feeding Behaviour	

Application Details

Application Notes:	Western blotting: 0.5-2 μg/mL	
	Immunohistochemistry: 5-20 μg/mL	
	Immunocytochemistry: 5-20 μg/mL	
	Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	

Handling

Format:	Liquid	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	24 months	



Western Blotting

Image 1. Detection of Recombinant OXTR, Rat using Polyclonal Antibody to Oxytocin Receptor (OXTR)