antibodies -online.com







anti-PDGFC antibody (AA 235-345)

2 Images



Go to Product page

\sim			
	N/6	1//r	$I \cap V$

Quantity:	100 μL	
Target:	PDGFC	
Binding Specificity:	AA 235-345	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PDGFC antibody is un-conjugated	
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

Product Details


Purpose:	Polyclonal Antibody to Platelet Derived Growth Factor C (PDGFC)	
Immunogen:	Recombinant Platelet Derived Growth Factor C (PDGFC) corresdonding to Val235~Gly345 (Accession # Q9NRA1)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against PDGFC. It has been selected for its ability to recognize PDGFC in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Pig, Rat	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	

Target Details

Expiry Date:


Target Details		
Target:	PDGFC Platelet Derived Growth Factor C (PDGFC Products) Viral Protein	
Alternative Name:		
Target Type:		
Background:	PDGF-C, SCDGF, VEGF-E, Fallotein, Spinal cord-derived growth factor, Platelet-derived growth factor C, latent form, Platelet-derived growth factor C, receptor-binding form	
Pathways:	RTK Signaling, Platelet-derived growth Factor Receptor Signaling	
Application Details		
Application Notes:	Western blotting: 0.5-2 μg/mL	
	1:250-1000 Immunohistochemistry: 5-20 μg/mL	
	1:25-100 Immunocytochemistry: 5-20 μg/mL	
	1:25-100 Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	
	detectable loss of activity. Avoid repeated freeze-thaw cycles.	
E . B .		

24 months



Immunohistochemistry

Image 1. Detection of PDGFC in Human Kidney Tissue using Polyclonal Antibody to Platelet Derived Growth Factor C (PDGFC)



Western Blotting

Image 2. Detection of Recombinant PDGFC, Human using Polyclonal Antibody to Platelet Derived Growth Factor C (PDGFC)