

Datasheet for ABIN7435547
anti-MEK1 antibody (AA 54-369)

3 Images

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	MEK1 (MAP2K1)
Binding Specificity:	AA 54-369
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MEK1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Mitogen Activated Protein Kinase Kinase 1 (MAP2K1)
Immunogen:	Recombinant Mitogen Activated Protein Kinase Kinase 1 (MAP2K1) corresponding to Leu54~Val369 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against MAP2K1. It has been selected for its ability to recognize MAP2K1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Pig, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details


Target:	MEK1 (MAP2K1)
Alternative Name:	Mitogen Activated Protein Kinase Kinase 1 (MAP2K1 Products)
Background:	MAPKK1, MEK1, MKK1, PRKMK1, Dual specificity mitogen-activated protein kinase kinase 1, ERK activator kinase 1
Pathways:	MAPK Signaling , RTK Signaling , Interferon-gamma Pathway , Fc-epsilon Receptor Signaling Pathway , Neurotrophin Signaling Pathway , Activation of Innate immune Response , Toll-Like Receptors Cascades , Autophagy , Signaling of Hepatocyte Growth Factor Receptor , BCR Signaling

Application Details

Application Notes:	Western blotting: 0.5-2 µg/mL Immunohistochemistry: 5-20 µg/mL Immunocytochemistry: 5-20 µg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months




Immunohistochemistry

Image 1. Detection of MAP2K1 in Human Cerebrum Tissue using Polyclonal Antibody to Mitogen Activated Protein Kinase Kinase 1 (MAP2K1)



Western Blotting

Image 2. Detection of Recombinant MAP2K1, Human using Polyclonal Antibody to Mitogen Activated Protein Kinase Kinase 1 (MAP2K1)



Immunohistochemistry

Image 3. Detection of MAP2K1 in Human Glioma Tissue using Polyclonal Antibody to Mitogen Activated Protein Kinase Kinase 1 (MAP2K1)