antibodies -online.com





anti-HMGCR antibody (AA 638-887)



Images



Go to Product page

\sim				
	$ V \cap$	r\/I	19	٨

Quantity:	100 μL
Target:	HMGCR
Binding Specificity:	AA 638-887
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HMGCR antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)


Product Details

Purpose:	Polyclonal Antibody to 3-Hydroxy-3-Methylglutaryl Coenzyme A Reductase (HMGCR)
Immunogen:	Recombinant 3-Hydroxy-3-Methylglutaryl Coenzyme A Reductase (HMGCR) corresdonding to Ala638~Ala887 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against HMGCR. It has been selected for its ability to recognize HMGCR in immunohistochemical staining and western blotting.
Cross-Reactivity:	Human, Mouse
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details


Target:	HMGCR	
Alternative Name:	3-Hydroxy-3-Methylglutaryl Coenzyme A Reductase (HMGCR Products)	
Background:	HMG-CoA Reductase, Hydroxymethylglutaryl-CoA Reductase	
Pathways:	AMPK Signaling, Negative Regulation of Hormone Secretion, Regulation of Lipid Metabolism by PPARalpha	

Pathways:	AMPK Signaling, Negative Regulation of Hormone Secretion, Regulation of Lipid Metabolism by PPARalpha
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL
	Immunohistochemistry: 5-20 μg/mL
	Immunocytochemistry: 5-20 μg/mL
	Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without
	detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months




Immunohistochemistry

Image 1. Detection of HMGCR in Rat Colon Tissue using
Polyclonal Antibody to 3-Hydroxy-3-Methylglutaryl
Coenzyme A Reductase (HMGCR)



Western Blotting

Image 2. Detection of Recombinant HMGCR, Rat using Polyclonal Antibody to 3-Hydroxy-3-Methylglutaryl Coenzyme A Reductase (HMGCR)



Immunohistochemistry

Image 3. Detection of HMGCR in Rat Cardiac Muscle Tissue using Polyclonal Antibody to 3-Hydroxy-3-Methylglutaryl Coenzyme A Reductase (HMGCR)

Please check the product details page for more images. Overall 7 images are available for ABIN7435559.