



[Go to Product page](#)

Datasheet for ABIN7435799
anti-CADM1 antibody (AA 27-292)

2 Images

Overview

Quantity:	100 µL
Target:	CADM1
Binding Specificity:	AA 27-292
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CADM1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Cell Adhesion Molecule 1 (CADM1)
Immunogen:	Recombinant Cell Adhesion Molecule 1 (CADM1) corresponding to Arg27~Val292 with N-terminal His and GST Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CADM1. It has been selected for its ability to recognize CADM1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Pig
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details


Target:	CADM1
Alternative Name:	Cell Adhesion Molecule 1 (CADM1 Products)
Background:	BL2, IGSF4, IGSF4A, NECL2, RA175, ST17, SYNCAM, TSLC1, sTSLC-1, synCAM1, Tumor Suppressor In Lung Cancer 1, Immunoglobulin Superfamily, Member 4, Nectin-Like 2
Pathways:	Regulation of Leukocyte Mediated Immunity , Positive Regulation of Immune Effector Process , Cell-Cell Junction Organization , Activated T Cell Proliferation

Application Details

Application Notes:	Western blotting: 1-5 µg/mL Immunocytochemistry in formalin fixed cells: 5-20 µg/mL Immunohistochemistry in formalin fixed frozen section: 5-20 µg/mL Immunohistochemistry in paraffin section: 5-20 µg/mL Enzyme-linked Immunosorbent Assay: 0.05-2 µg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



Western Blotting

Image 1. Detection of CADM1 in Porcine Cerebrum lysate using Polyclonal Antibody to Cell Adhesion Molecule 1 (CADM1)

Western Blotting

Image 2. Detection of Recombinant CADM1, Human using Polyclonal Antibody to Cell Adhesion Molecule 1 (CADM1)