

Datasheet for ABIN7435910  
**anti-HDAC9 antibody (AA 23-343)**

## 3 Images

[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	HDAC9
Binding Specificity:	AA 23-343
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HDAC9 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

## Product Details

Purpose:	Polyclonal Antibody to Histone Deacetylase 9 (HDAC9)
Immunogen:	Recombinant Histone Deacetylase 9 (HDAC9) corresponding to Pro23~Ala343
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against HDAC9. It has been selected for its ability to recognize HDAC9 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

## Target Details


Target:	HDAC9
Alternative Name:	Histone Deacetylase 9 ( <a href="#">HDAC9 Products</a> )
Background:	HD7, HDAC, HDAC7, HDAC7B, HDAC9B, HDAC9FL, HDRP, MITR, Histone deacetylase-related protein, MEF2-interacting transcription repressor MITR, Histone deacetylase 7B
Pathways:	<a href="#">Regulation of Muscle Cell Differentiation</a> , <a href="#">Skeletal Muscle Fiber Development</a>

## Application Details

Application Notes:	Western blotting: 0.5-2 µg/mL Immunohistochemistry: 5-20 µg/mL Immunocytochemistry: 5-20 µg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



#### Western Blotting

**Image 1.** Detection of HDAC9 in Human K562 cell lysate using Polyclonal Antibody to Histone Deacetylase 9 (HDAC9)

#### Western Blotting

**Image 2.** Detection of Recombinant HDAC9, Human using Polyclonal Antibody to Histone Deacetylase 9 (HDAC9)

#### Western Blotting

**Image 3.** Detection of HDAC9 in Human K562 cell lysate using Polyclonal Antibody to Histone Deacetylase 9 (HDAC9)