# antibodies - online.com







# anti-SIRT1 antibody (AA 236-490)

**Images** 



# Overview

Quantity:	100 μL
Target:	SIRT1
Binding Specificity:	AA 236-490
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SIRT1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

# **Product Details**

Durnoon	Dalvalanal Antibady to Cirtuin 1 (CIDT1)
Purpose:	Polyclonal Antibody to Sirtuin 1 (SIRT1)
Immunogen:	Recombinant Sirtuin 1 (SIRT1) corresdonding to Glu236~Ala490 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against SIRT1. It has been selected for its
	ability to recognize SIRT1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Human, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

# **Target Details**

- Target Betane	
Target:	SIRT1
Alternative Name:	Sirtuin 1 (SIRT1 Products)
Background:	SIR2L1, Silent Mating Type Information Regulation 2 Homolog 1, NAD-Dependent Deacetylase Sirtuin-1, SIR2-like protein 1, SirtT1 75 kDa fragment
Pathways:	MAPK Signaling, Intracellular Steroid Hormone Receptor Signaling Pathway, Regulation of Intracellular Steroid Hormone Receptor Signaling, Carbohydrate Homeostasis, Positive Regulation of Endopeptidase Activity, Regulation of Carbohydrate Metabolic Process, Positive Regulation of Response to DNA Damage Stimulus, Negative Regulation of intrinsic apoptotic Signaling
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-20 μg/mL Optimal working dilutions must be determined by end user.
	-p

### Comment:


The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### Restrictions:

For Research Use only


# Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



# **Immunohistochemistry**

Image 1. #VALUE!



# **Western Blotting**

Image 2. Detection of Recombinant SIRT1, Mouse using Polyclonal Antibody to Sirtuin 1 (SIRT1)