# antibodies - online.com







## anti-HSD17B3 antibody (AA 1-310)

**Images** 



#### Overview


Quantity:	100 μL
Target:	HSD17B3
Binding Specificity:	AA 1-310
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HSD17B3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

#### **Product Details**

Purpose:	Polyclonal Antibody to 17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3)
Immunogen:	Recombinant 17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3) corresdonding to Met1~Arg310 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against HSD17b3. It has been selected for its ability to recognize HSD17b3 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Rabbit
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography


### **Target Details**

Target Details	
Target:	HSD17B3
Alternative Name:	17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17B3 Products)
Background:	SDR12C2, EDH17B3, Short Chain Dehydrogenase/Reductase Family 12C,Member 2,
	Testosterone 17-Beta-Dehydrogenase 3, Testicular 17-beta-hydroxysteroid dehydrogenase
Pathways:	Metabolism of Steroid Hormones and Vitamin D, Steroid Hormone Biosynthesis
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL
	Immunohistochemistry: 5-20 μg/mL
	Immunocytochemistry: 5-20 μg/mL
	Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without
	detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



#### **Immunohistochemistry**

Image 1. #VALUE!



#### **Western Blotting**

Image 2. Detection of Recombinant HSD17b3, Human usingPolyclonal Antibody to 17-Beta-HydroxysteroidDehydrogenase Type 3 (HSD17b3)