antibodies -online.com





anti-PDE10A antibody (AA 508-779)

2 Images



Go to Product page

Overview

Quantity:	100 μL
Target:	PDE10A
Binding Specificity:	AA 508-779
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PDE10A antibody is un-conjugated
Application:	Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (IHC), Immunocytochemistry (ICC)


Product Details

Purpose:	Polyclonal Antibody to Phosphodiesterase 10A (PDE10A)
Immunogen:	Recombinant Phosphodiesterase 10A (PDE10A) corresdonding to Asn508~Asp779 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against PDE10A. It has been selected for its ability to recognize PDE10A in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details


Target:	PDE10A
Alternative Name:	Phosphodiesterase 10A (PDE10A Products)
Background:	CAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A
Pathways:	cAMP Metabolic Process

Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL
	Immunohistochemistry: 5-20 μg/mL
	Immunocytochemistry: 5-20 μg/mL
	Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide



Immunohistochemistry

Image 1. #VALUE!



Western Blotting

Image 2. Detection of Recombinant PDE10A, Human using Polyclonal Antibody to Phosphodiesterase 10A (PDE10A)