antibodies -online.com

anti-Syncoilin antibody (AA 169-458)









Go to Product page

Overview

Quantity:	100 μL
Target:	Syncoilin (Sync)
Binding Specificity:	AA 169-458
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Syncoilin antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details


Purpose:	Polyclonal Antibody to Syncoilin (SYNC)
Immunogen:	Recombinant Syncoilin (SYNC) corresdonding to Ile169~Pro458 with N-terminal His and GST Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against SYNC. It has been selected for its ability to recognize SYNC in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Pig, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Expiry Date:

Target Details	
Target:	Syncoilin (Sync)
Alternative Name:	Syncoilin (Sync Products)
Background:	Syncoilin-1, Intermediate Filament Protein, Syncoilin intermediate filament 1
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL
	Immunohistochemistry: 5-20 μg/mL
	Immunocytochemistry: 5-20 μg/mL
	Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without

detectable loss of activity. Avoid repeated freeze-thaw cycles.

24 months



Western Blotting

Image 1. Detection of Recombinant SYNC, Human using Polyclonal Antibody to Syncoilin (SYNC)