

Datasheet for ABIN7436183  
**anti-Stanniocalcin 2 antibody (AA 53-294)**[Go to Product page](#)

## 3 Images

## Overview

Quantity:	100 µL
Target:	Stanniocalcin 2 (STC2)
Binding Specificity:	AA 53-294
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Stanniocalcin 2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

## Product Details

Purpose:	Polyclonal Antibody to Stanniocalcin 2 (STC2)
Immunogen:	Recombinant Stanniocalcin 2 (STC2) corresponding to Ile53~Gln294 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against STC2. It has been selected for its ability to recognize STC2 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

## Target Details


Target:	Stanniocalcin 2 (STC2)
Alternative Name:	Stanniocalcin 2 ( <a href="#">STC2 Products</a> )
Background:	STCRP, Stanniocalcin-related protein
Pathways:	<a href="#">Hormone Activity</a> , <a href="#">ER-Nucleus Signaling</a> , <a href="#">SARS-CoV-2 Protein Interactome</a>

## Application Details

Application Notes:	Western blotting: 0.5-2 µg/mL Immunohistochemistry: 5-20 µg/mL Immunocytochemistry: 5-20 µg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



**Western Blotting**

**Image 1.** Detection of STC2 in Mouse Liver lysate using Polyclonal Antibody to Stanniocalcin 2 (STC2)

**Western Blotting**

**Image 2.** Detection of Recombinant STC2, Human using Polyclonal Antibody to Stanniocalcin 2 (STC2)

**Western Blotting**

**Image 3.** Detection of STC2 in Mouse Skeletal muscle lysate using Polyclonal Antibody to Stanniocalcin 2 (STC2)