antibodies .- online.com







anti-HDC antibody (AA 160-369)





\sim	
()\/\	view
	V I C V V

Quantity:	100 μL
Target:	HDC
Binding Specificity:	AA 160-369
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HDC antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Histidine Decarboxylase (HDC)
Immunogen:	Recombinant Histidine Decarboxylase (HDC) corresdonding to Ala160~Phe369 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against HDC. It has been selected for its ability to recognize HDC in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target: HDC


Target Details

Histidine Decarboxylase (HDC Products) Alternative Name:

Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL
	Immunohistochemistry: 5-20 μg/mL
	Immunocytochemistry: 5-20 μg/mL
	Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

Expiry Date: 24 months




Western Blotting

Image 1. Detection of HDC in Human 293T cell lysate using Polyclonal Antibody to Histidine Decarboxylase (HDC)



Western Blotting

Image 2. Detection of Recombinant HDC, Human using Polyclonal Antibody to Histidine Decarboxylase (HDC)



Immunohistochemistry

Image 3. #VALUE!