
antibodies -online.com





anti-PPARGC1A antibody (AA 573-767)





Go to Product page

| 1//(| | | |
|------|--|--|--|
| | | | |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | PPARGC1A | |
| Binding Specificity: | AA 573-767 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This PPARGC1A antibody is un-conjugated | |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP) | |
| Product Details | | |
| Purpose: | Polyclonal Antibody to Peroxisome Proliferator Activated Receptor Gamma Coactivator 1 Alpha (PPARgC1a) | |
| Immunogen: | Recombinant Peroxisome Proliferator Activated Receptor Gamma Coactivator 1 Alpha (PPARgC1a) corresdonding to Ser573~Asn767 with N-terminal His Tag | |
| | | |
| Isotype: | IgG | |
| Specificity: | IgG The antibody is a rabbit polyclonal antibody raised against PPARgC1a. It has been selected for its ability to recognize PPARgC1a in immunohistochemical staining and western blotting. | |
| | The antibody is a rabbit polyclonal antibody raised against PPARgC1a. It has been selected for | |

Target Details


| Target: | PPARGC1A |
|-------------------|---|
| Alternative Name: | Peroxisome Proliferator Activated Receptor Gamma Coactivator 1 Alpha (PPARGC1A Products) |
| Background: | LEM6, PGC-1(alpha), PGC-1v, PGC1, PGC1A, PPARGC1, Ligand effect modulator 6 |
| Pathways: | Intracellular Steroid Hormone Receptor Signaling Pathway, Carbohydrate Homeostasis, Regulation of Lipid Metabolism by PPARalpha, Regulation of Carbohydrate Metabolic Process, Smooth Muscle Cell Migration, Brown Fat Cell Differentiation |

Application Details

| Application Notes: | Western blotting: 0.5-2 µg/mL |
|--------------------|--|
| | Immunohistochemistry: 5-20 μg/mL |
| | Immunocytochemistry: 5-20 μg/mL |
| | Optimal working dilutions must be determined by end user. |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated |
| | thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious |
| | degradation and precipitation were observed. The loss rate is less than 5% within the expiration |
| | date under appropriate storage condition. |
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid |
|--------------------|---|
| Buffer: | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles. |
| Expiry Date: | 24 months |



Western Blotting

Image 1. Detection of Recombinant PPARgC1a, Human using Polyclonal Antibody to Peroxisome Proliferator Activated Receptor Gamma Coactivator 1 Alpha (PPARgC1a)