

Datasheet for ABIN7436513
anti-MPG antibody (AA 221-296)



[Go to Product page](#)

5 Images

Overview

Quantity:	100 µL
Target:	MPG
Binding Specificity:	AA 221-296
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MPG antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Purpose:	Polyclonal Antibody to N-Methylpurine DNA Glycosylase (MPG)
Immunogen:	Recombinant N-Methylpurine DNA Glycosylase (MPG) corresponding to Pro70~Lys220 and Leu221~Thr296 linked by a peptide LCQALA with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against MPG. It has been selected for its ability to recognize MPG in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details


Target:	MPG
Alternative Name:	N-Methylpurine DNA Glycosylase (MPG Products)
Background:	AAG, APNG, ADPG, CRA36.1, MDG, Mid1, PIG11, PIG16, Anpg, DNA-3-Methyladenine Glycosylase, Alkyladenine DNA Glycosylase, 3-alkyladenine DNA glycosylase
Pathways:	DNA Damage Repair

Application Details

Application Notes:	Western blotting: 1-5 µg/mL Immunocytochemistry in formalin fixed cells: 5-20 µg/mL Immunohistochemistry in formalin fixed frozen section: 5-20 µg/mL Immunohistochemistry in paraffin section: 5-20 µg/mL Enzyme-linked Immunosorbent Assay: 0.05-2 µg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months




Immunohistochemistry

Image 1. Detection of MPG in Human Rectum Tissue using Polyclonal Antibody to N-Methylpurine DNA Glycosylase (MPG)



Western Blotting

Image 2. Detection of Recombinant MPG, Human using Polyclonal Antibody to N-Methylpurine DNA Glycosylase (MPG)



Immunohistochemistry

Image 3. Detection of MPG in Human Breast cancer Tissue using Polyclonal Antibody to N-Methylpurine DNA Glycosylase (MPG)

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN7436513.