antibodies .- online.com







anti-MOGAT2 antibody (AA 48-217)



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	MOGAT2	
Binding Specificity:	AA 48-217	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This MOGAT2 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

Product Details

Purpose:	Polyclonal Antibody to Monoacylglycerol-O-acyltransferase 2 (MOGAT2)	
Immunogen:	Recombinant Monoacylglycerol-O-acyltransferase 2 (MOGAT2) corresdonding to Leu48~Gly217 with N-terminal His Tag	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against MOGAT2. It has been selected for its ability to recognize MOGAT2 in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	

Target Details

Target:	MOGAT2	
rarget.	WIOGATZ	

Target Details

g		
Alternative Name:	Monoacylglycerol-O-acyltransferase 2 (MOGAT2 Products)	
Background:	MGAT2, DGAT2L5, DC5, 2-Acylglycerol O-Acyltransferase 2, Diacylglycerol O-acyltransferase	
	candidate 5, Diacylglycerol acyltransferase 2-like protein 5	
Application Details		
Application Notes:	Western blotting: 0.5-2 μg/mL	
	Immunohistochemistry: 5-20 µg/mL	
	Immunocytochemistry: 5-20 μg/mL	
	Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	
	detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	24 months	