

Datasheet for ABIN7436747  
**anti-PIK3CD antibody (AA 774-1044)**[Go to Product page](#)

## 3 Images

## Overview

Quantity:	100 µL
Target:	PIK3CD
Binding Specificity:	AA 774-1044
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PIK3CD antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

## Product Details

Purpose:	Polyclonal Antibody to Phosphoinositide-3-Kinase Catalytic Delta Polypeptide (PIK3Cd)
Immunogen:	Recombinant Phosphoinositide-3-Kinase Catalytic Delta Polypeptide (PIK3Cd) corresponding to Val774~Gln1044 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against PIK3Cd. It has been selected for its ability to recognize PIK3Cd in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

## Target Details


Target:	PIK3CD
Alternative Name:	Phosphoinositide-3-Kinase Catalytic Delta Polypeptide ( <a href="#">PIK3CD Products</a> )
Background:	P110D, P110δ, Phosphoinositide-3-Kinase C, Phosphatidylinositol 4,5-bisphosphate 3-kinase 110 kDa catalytic subunit delta
Pathways:	<a href="#">BCR Signaling</a> , <a href="#">Warburg Effect</a>

## Application Details

Application Notes:	Western blotting: 1-5 µg/mL Immunocytochemistry in formalin fixed cells: 5-20 µg/mL Immunohistochemistry in formalin fixed frozen section: 5-20 µg/mL Immunohistochemistry in paraffin section: 5-20 µg/mL Enzyme-linked Immunosorbent Assay: 0.05-2 µg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months




### Immunohistochemistry

**Image 1.** Detection of PIK3Cd in Human Colorectal cancer Tissue using Polyclonal Antibody to Phosphoinositide-3-Kinase Catalytic Delta Polypeptide (PIK3Cd)



### Western Blotting

**Image 2.** Detection of PIK3Cd in Mouse Liver Tissue using Polyclonal Antibody to Phosphoinositide-3-Kinase Catalytic Delta Polypeptide (PIK3Cd)



### Western Blotting

**Image 3.** Detection of Recombinant PIK3Cd, Human using Polyclonal Antibody to Phosphoinositide-3-Kinase Catalytic Delta Polypeptide (PIK3Cd)