# antibodies -online.com





## anti-PRKDC antibody (AA 3899-4128)

2 Images



Go to Product page

$\sim$	
( )\/△	rview
$\cup$	1 410 44

Quantity:	100 μL
Target:	PRKDC
Binding Specificity:	AA 3899-4128
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PRKDC antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

#### **Product Details**

Purpose:	Polyclonal Antibody to Protein Kinase, DNA Activated, Catalytic Polypeptide (PRKDC)
Immunogen:	Recombinant Protein Kinase, DNA Activated, Catalytic Polypeptide (PRKDC) corresdonding to Ala3899~Met4128
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against PRKDC. It has been selected for its ability to recognize PRKDC in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

### **Target Details**

Target:	PRKDC	
rarget.	TINDO	

## **Target Details**


Alternative Name:	Protein Kinase, DNA Activated, Catalytic Polypeptide (PRKDC Products)
Background:	HYRC, HYRC1, DNPK1, p350, DNAPK, XRCC7, DNA-PKcs
Pathways:	DNA Damage Repair, Production of Molecular Mediator of Immune Response

## **Application Details**

Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL
	1:400-1600 Immunohistochemistry: 5-20 μg/mL
	1:40-160 Immunocytochemistry: 5-20 µg/mL
	1:40-160 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	


## Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



#### **Immunohistochemistry**

**Image 1.** Detection of PRKDC in Mouse Uterus Tissue using Polyclonal Antibody to Protein Kinase, DNA Activated, Catalytic Polypeptide (PRKDC)



#### **Western Blotting**

Image 2. Detection of Recombinant PRKDC, Mouse using Polyclonal Antibody to Protein Kinase, DNA Activated, Catalytic Polypeptide (PRKDC)