

Datasheet for ABIN7437071 anti-TNF alpha antibody (AA 80-235) (APC)

2 Images




Go to Product page

\sim			
	ve	r\/	٨
\cup	V C	1 V I	٧V

Quantity:	100 μL	
Target:	TNF alpha	
Binding Specificity:	AA 80-235	
Reactivity:	Rabbit	
Host:	Guinea Pig	
Clonality:	Polyclonal	
Conjugate:	This TNF alpha antibody is conjugated to APC	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	
Product Details	Delivelenal Antibady to Tumor Negrocia Factor Alpha (TNFa)	
Purpose:	Polyclonal Antibody to Tumor Necrosis Factor Alpha (TNFa)	
Immunogen:	Recombinant Tumor Necrosis Factor Alpha (TNFa) corresdonding to Leu80~Leu235 (Accession # P04924)	
Isotype:	IgG	
Specificity:	The antibody is a cavia polyclonal antibody raised against TNFa. It has been selected for its ability to recognize TNFa in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	
Target Details		
Target:	TNF alpha	


Target Details

Alternative Name:	TNFa (TNF alpha Products)	
Background:	DIF, TNF-A, TNFSF2, Cachectin, Tumor Necrosis Factor Ligand Superfamily Member 2	
UniProt:	P04924	
Pathways:	NF-kappaB Signaling, Apoptosis, Caspase Cascade in Apoptosis, TLR Signaling, Cellular	
	Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positive	
	Regulation of Immune Effector Process, Production of Molecular Mediator of Immune	
	Response, Positive Regulation of Endopeptidase Activity, Hepatitis C, Protein targeting to	
	Nucleus, Inflammasome	
Application Details		
Application Notes:	Western blotting: 0.5-2 μg/mL,1:500-2000,Immunohistochemistry: 5-20 μg/mL,1:50-	
	200,lmmunocytochemistry: 5-20 μg/mL,1:50-200,Optimal working dilutions must be	
	determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	500 μg/mL	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be	
	handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	
	detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	24 months	
Lapity Date.	Z# HUUHUIO	



Immunohistochemistry

Image 1. Detection of TNFa in Rabbit Cerebrum Tissue using Polyclonal Antibody to Tumor Necrosis Factor Alpha (TNFa)



Western Blotting

Image 2. Detection of Recombinant TNFa, Rabbit using Polyclonal Antibody to Tumor Necrosis Factor Alpha (TNFa)