antibodies -online.com





anti-ICOS antibody (AA 35-131)



Image



Go to Product page

Overview

Quantity:	100 μL
Target:	ICOS
Binding Specificity:	AA 35-131
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ICOS antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Inducible T-Cell Co Stimulator (ICOS)
Immunogen:	Recombinant Inducible T-Cell Co Stimulator (ICOS) corresdonding to Asn35~Ile131 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against ICOS. It has been selected for its ability to recognize ICOS in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography


Target Details

Target Details

Alternative Name:	Inducible T-Cell Co Stimulator (ICOS Products)
Background:	CD278, AILIM, Inducible T-Cell Costimulator, Activation-Inducible Lymphocyte Immunomediatory Molecule
Pathways:	Cancer Immune Checkpoints

Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL
	1:500-2000 Immunohistochemistry: 5-20 μg/mL
	1:50-200 Immunocytochemistry: 5-20 µg/mL
	1:50-200 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Format:	Liquid
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



Western Blotting

Image 1. Detection of Recombinant ICOS, Mouse using Polyclonal Antibody to Inducible T-Cell Co Stimulator (ICOS)