antibodies -online.com





anti-TPT1 antibody (AA 1-172)

2 Images



Go to Product page

Overview

| Quantity: | 100 μL |
|----------------------|--|
| Target: | TPT1 |
| Binding Specificity: | AA 1-172 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This TPT1 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

Product Details


| Purpose: | Polyclonal Antibody to Tumor Protein, Translationally Controlled 1 (TPT1) |
|-------------------|--|
| Immunogen: | Recombinant Tumor Protein, Translationally Controlled 1 (TPT1) corresdonding to Met1~Cys172 (Accession # P13693) |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against TPT1. It has been selected for its ability to recognize TPT1 in immunohistochemical staining and western blotting. |
| Cross-Reactivity: | Mouse, Rat |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography |

Target Details

Expiry Date:


24 months

| rarget Details | |
|---------------------|---|
| Target: | TPT1 |
| Alternative Name: | Tumor Protein, Translationally Controlled 1 (TPT1 Products) |
| Background: | HRF, TCTP, p02, p23, Fortilin, Translationally-Controlled Tumor Protein, Histamine-releasing factor |
| Pathways: | Response to Water Deprivation, Regulation of Cell Size, Negative Regulation of intrinsic apoptotic Signaling |
| Application Details | |
| Application Notes: | Western blotting: 0.5-2 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-20 μg/mL Optimal working dilutions must be determined by end user. |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles. |



Immunohistochemistry

Image 1. #VALUE!



Western Blotting

Image 2. Detection of Recombinant TPT1, Human using Polyclonal Antibody to Tumor Protein, Translationally Controlled 1 (TPT1)