antibodies .- online.com





anti-PPARGC1A antibody (AA 577-766)

IgG

Human, Rat





Go to Product page

1//(

Isotype:

Specificity:

Cross-Reactivity:

Purification:

Quantity:	100 μL		
Target:	PPARGC1A		
Binding Specificity:	AA 577-766		
Reactivity:	Mouse		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This PPARGC1A antibody is un-conjugated		
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)		
Product Details			
Purpose:	Polyclonal Antibody to Peroxisome Proliferator Activated Receptor Gamma Coactivator 1 Alpha (PPARgC1a)		
Immunogen:	Recombinant Peroxisome Proliferator Activated Receptor Gamma Coactivator 1 Alpha (PPARgC1a) corresdonding to Cys577~Asn766 with N-terminal His Tag		

The antibody is a rabbit polyclonal antibody raised against PPARgC1a. It has been selected for

its ability to recognize PPARgC1a in immunohistochemical staining and western blotting.

Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details


Target:	PPARGC1A	
Alternative Name:	Peroxisome Proliferator Activated Receptor Gamma Coactivator 1 Alpha (PPARGC1A Products)	
Background:	LEM6, PGC-1(alpha), PGC-1v, PGC1, PGC1A, PPARGC1, Ligand effect modulator 6	
Pathways:	Intracellular Steroid Hormone Receptor Signaling Pathway, Carbohydrate Homeostasis, Regulation of Lipid Metabolism by PPARalpha, Regulation of Carbohydrate Metabolic Process, Smooth Muscle Cell Migration, Brown Fat Cell Differentiation	

Application Details

Application Notes:	Western blotting: 0.5-2 µg/mL		
	Immunohistochemistry: 5-20 μg/mL		
	Immunocytochemistry: 5-20 μg/mL		
	Optimal working dilutions must be determined by end user.		
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated		
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious		
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration		
	date under appropriate storage condition.		
Restrictions:	For Research Use only		

Handling

Tariding		
Format:	Liquid	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	24 months	



Western Blotting

Image 1. Detection of Recombinant PPARgC1a, Mouse using Polyclonal Antibody to Peroxisome Proliferator Activated Receptor Gamma Coactivator 1 Alpha (PPARgC1a)