

Datasheet for ABIN7437863  
**anti-Acrosin antibody (AA 43-291)**[Go to Product page](#)

## 2 Images

## Overview

|                      |  |
|----------------------|--|
| Quantity:            | 100 µL   |
| Target:              | Acrosin (ACR)  |
| Binding Specificity: | AA 43-291  |
| Reactivity:          | Rat  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This Acrosin antibody is un-conjugated   |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP) |

## Product Details

|                   |  |
|-------------------|--|
| Purpose:          | Polyclonal Antibody to Acrosin (ACR)   |
| Immunogen:        | Recombinant Acrosin (ACR) corresponding to Ile43~Gly291  |
| Isotype:          | IgG  |
| Specificity:      | The antibody is a rabbit polyclonal antibody raised against ACR. It has been selected for its ability to recognize ACR in immunohistochemical staining and western blotting. |
| Cross-Reactivity: | Mouse  |
| Purification:     | Antigen-specific affinity chromatography followed by Protein A affinity chromatography   |

## Target Details


|                   |  |
|-------------------|--|
| Target:           | Acrosin (ACR)                            |
| Alternative Name: | Acrosin ( <a href="#">ACR Products</a> ) |
| Background:       | ACRS, Preproacrosin                      |
| Pathways:         | <a href="#">cAMP Metabolic Process</a>   |

## Application Details

|                    |   |
|--------------------|---|
| Application Notes: | Western blotting: 0.5-2 µg/mL Immunohistochemistry: 5-20 µg/mL Immunocytochemistry: 5-20 µg/mL Optimal working dilutions must be determined by end user.  |
| Comment:           | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions:      | For Research Use only   |


## Handling

|                    |   |
|--------------------|---|
| Format:            | Liquid  |
| Buffer:            | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.  |
| Preservative:      | ProClin   |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.   |
| Storage:           | 4 °C,-20 °C   |
| Storage Comment:   | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles. |
| Expiry Date:       | 24 months   |



Immunohistochemistry

**Image 1.** Detection of ACR in Rat Intestine Tissue using Polyclonal Antibody to Acrosin (ACR)



Western Blotting

**Image 2.** Detection of Recombinant ACR, Rat using Polyclonal Antibody to Acrosin (ACR)