# antibodies -online.com





# anti-PLAUR antibody (AA 221-305)

2 Images



Go to Product page

$\sim$	
( )\/△	rview
$\cup$	1 410 44

Quantity:	100 μL
Target:	PLAUR
Binding Specificity:	AA 221-305
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PLAUR antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

### **Product Details**

Purpose:	Polyclonal Antibody to Plasminogen Activator, Urokinase Receptor (uPAR)
Immunogen:	Recombinant Plasminogen Activator, Urokinase Receptor (uPAR) corresdonding to Gly221~Gly305 (Accession # Q03405) with N-terminal His and GST Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against uPAR. It has been selected for its ability to recognize uPAR in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

## **Target Details**

Target: PLAUR
---------------

## **Target Details**


Alternative Name:	Plasminogen Activator, Urokinase Receptor (PLAUR Products)
Background:	CD87, PLAUR, PLAU-R, U-PAR, URKR, Monocyte Activation Antigen Mo3
Pathways:	Inositol Metabolic Process

## **Application Details**

Application Notes:	Western blotting: 0.5-2 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-
	20 μg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



### **Immunohistochemistry**

Image 1. #VALUE!



### **Western Blotting**

**Image 2.** Detection of Recombinant uPAR, Human using Polyclonal Antibody to Plasminogen Activator, Urokinase Receptor (uPAR)